

Sensibilitat *Mycobacterium tuberculosis* 2002-2020 HSCSP

2019-2020										
Antecedents de tractament		Sense tractament previ			Amb tractament anterior			Antecedent de tractament desconeugut		
		total	autòcton	origen estranger	total	autòcton	origen estranger	total	autòcton	origen estranger
sensibles	44	38	18	20	3	2	1	3		3
resistents	38 (86.4%)	32 (84.2%)	15 (83.3%)	17 (85%)	3 (100)	2	1	3 (100)		3
Isoniazida	6 (13.6%)	6 (15.8%)	3 (16.7%)	3 (15%)						
Rifampicina	1(2.3%)		1 (5%)							
Etambutol	1(2.3%)		1 (5.5%)							
Estreptomicina	4 (9.1%)		2 (11.1%)	2 (10%)						
Pirazinamida										
:										
MDR										

2002-2020										
Antecedents de tractament	TOTAL	Sense tractament previ			Amb tractament anterior			Antecedent de tractament desconeugut		
		total	autòcton	origen estranger	total	autòcton	origen estranger	total	autòcton	origen estranger
sensibles	757	580	380	200	70	44	26	107	73	34
resistents	670 (88.5%)	512 (88.3%)	355 (93.4%)	157 (78.5%)	59 (84.3%)	40 (90.9%)	19 (73.1%)	99 (92,5%)	68 (93,2%)	31 (91,2%)
	87 (11.5%)	68 (11.7%)	25 (6.6%)	43 (21.5%)	11 (15,7%)	4 (9,1%)	7 (26,9%)	8 (7,5%)	5 (6,8%)	3 (8,8%)
Isoniazida	51 (6.7%)	39 (6,7%)	17 (4,5%)	23 (11,5%)	7 (10%)	2 (4,5%)	5 (19,2%)	5 (4,7%)	3 (4,1%)	2 (5,9%)
Rifampicina	13 (1.7%)	9 (1,6%)	0	9 (4,5%)	4 (5,7%)	2 (4,5)	2 (7,7%)	0	0	0
Etambutol	7 (0.9%)	5 (1%)	3 (0,8%)	3 (1,5%)	2 (2,9%)	0	2 (7,7%)	0	0	0
Estreptomicina	23 (3.0%)	17 (2,9%)	7 (1,8%)	14 (7%)	6 (8,5%)	3 (6,8%)	3 (11,5%)	4 (3,7%)	3 (4,1%)	1 (2,9%)
Pirazinamida	16 (2.1%)	11 (1,9%)	3 (0,8%)	9 (4,5%)	5 (7,1%)	2 (4,5%)	3 (11,5%)	0	0	0
MDR	9 (1.2%)	6 (1,03%)	0	6 (3%)	4(5,7%)	2 (4,5%)	2 (7,7%)	0	0	0